

One-sided Limits and Continuity

One-Sided Limits:

Right-Hand Limit: $\lim_{x \rightarrow a^+} f(x) = L$ means that $f(x) \rightarrow L$ as $x \rightarrow a$
from the right ($x > a$)

Left-Hand Limit: $\lim_{x \rightarrow a^-} f(x) = L$ means that $f(x) \rightarrow L$ as $x \rightarrow a$
from the left ($x < a$)

Theorem: The ordinary limit $\lim_{x \rightarrow a} f(x)$ exists if and only if both corresponding one-sided limits exist and are equal.

$$\lim_{x \rightarrow a} f(x) = L \Leftrightarrow \lim_{x \rightarrow a^+} f(x) = L \Leftrightarrow \lim_{x \rightarrow a^-} f(x) = L$$

Example: $f(x) = \sqrt{x - 4}$

$$\lim_{x \rightarrow 4^+} \sqrt{x - 4} =$$

$$\lim_{x \rightarrow 4^-} \sqrt{x - 4} =$$

$$\lim_{x \rightarrow 4} \sqrt{x - 4} =$$

All limit theorems hold.

$$\lim_{x \rightarrow 2^+} \frac{2 - x}{x^2 - 2}$$

$$\lim_{x \rightarrow 4^-} \frac{\sqrt{x} - 2}{x - 4}$$

$$\lim_{x \rightarrow 4^-} \frac{|x - 4|}{x - 4}$$

$$\lim_{x \rightarrow 1^+} \frac{|x - 1|}{1 - x}$$

$$\lim_{x \rightarrow 0^-} \lceil \lceil x \rceil \rceil$$

$$\lim_{x \rightarrow 1^+} \lceil \lceil x + 2 \rceil \rceil$$

$$\lim_{x \rightarrow 2^+} (2x - \lceil x \rceil)$$

$$\lim_{x \rightarrow 1} f(x) = \begin{cases} 3x - 1 & , \quad x < 1 \\ x^2 & , \quad x > 1 \end{cases}$$

$$\lim_{x \rightarrow 1^-} f(x) =$$

$$\lim_{x \rightarrow 1^+} f(x) =$$

$$\lim_{x \rightarrow 1} f(x) =$$

$$\lim_{x \rightarrow 2} \text{if } f(x) = \begin{cases} x^2 & , \quad x < 2 \\ 3x - 2 & , \quad x \geq 2 \end{cases}$$




$$\lim_{x \rightarrow 2^-} f(x) =$$

$$\lim_{x \rightarrow 2^+} f(x) =$$

$$\lim_{x \rightarrow 2} f(x) =$$

Continuity:

A function is continuous at $x = c$ if it has no holes, jumps or gaps at c .



Continuity at a Point: A function f is continuous at a point c if the following three conditions exist:

- 1.
- 2.
- 3.

Continuity on an Interval: A function f is continuous on an open interval (a, b) if it is continuous at each point in the interval. A function that is continuous on the entire real line $(-\infty, +\infty)$ is said to be "everywhere" continuous.





Consider a function such as:

What would you say about the continuity of this function?

A function f is continuous on the closed interval $[a, b]$ if it is continuous at each point on the open interval (a, b) and if $\lim_{x \rightarrow a^+} f(x) = f(a)$ (f is continuous from the right at a) and $\lim_{x \rightarrow b^-} f(x) = f(b)$ (f is continuous from the left at b).

There are two classifications of a point of discontinuity on a function.

1. Removable: if $\lim_{x \rightarrow a} f(x)$ exists; function can be redefined at a to remove the discontinuity.
2. Non-removable: if $\lim_{x \rightarrow a} f(x)$ does not exist.



Properties of Continuity

If b is a real number and f and g are continuous at $x = c$, then the following functions are also continuous at c .

1. A scalar multiple: bf
2. A sum or difference: $f \pm g$
3. A product: fg
4. A quotient: $\frac{f}{g}$

The following types of functions are continuous at every point in their domains.

1. Polynomial
2. Rational
3. Radical
4. Trigonometric

Find x – values (if any) at which f is not continuous. Classify each discontinuity as removable or non-removable. If removable, redefine the function to remove the discontinuity.

$$f(x) = \frac{1}{x^2 + 1}$$

$$f(x) = \frac{2}{x - 1}$$

$$f(x) = \frac{x - 3}{x^2 - 9}$$


$$f(x) = \frac{|x - 3|}{x - 3}$$

$$f(x) = \begin{cases} -2x & \text{if } x \leq 2 \\ x^2 - 4x + 1 & \text{if } x > 2 \end{cases}$$


$$f(x) = \begin{cases} \tan \frac{\pi x}{4} & \text{if } |x| < 1 \\ x & \text{if } |x| \geq 1 \end{cases}$$

State the intervals of continuity:

58. $f(x) = x\sqrt{x+3}$



60. $f(x) = \frac{x+1}{\sqrt{x}}$




Find "a" so that f is continuous for all real numbers x .


$$f(x) = \begin{cases} \frac{x^2 - a^2}{x - a} & \text{if } x \neq a \\ 8 & \text{if } x = a \end{cases}$$

Intermediate Value Theorem

If f is continuous on the closed interval $[a, b]$ and k is any number between $f(a)$ and $f(b)$, then there is at least one number c in $[a, b]$ such that $f(c) = k$.



f is continuous on $[a, b]$.
(For k , there exist 3 c 's.)



f is not continuous on $[a, b]$.
(For k , there are no c 's.)

A continuous function on $[a, b]$ takes on all values between $f(a)$ and $f(b)$.

Use the Intermediate Value Theorem to show that $f(x) = x^3 + 2x - 1$ has a zero in the interval $[0, 1]$.

Use the Intermediate Value Theorem to prove that $\sqrt{3}$ is between 1 and 2.