

Dividing Polynomials Video Lecture

Section 4.2

Course Learning Objectives:

Demonstrate appropriate manipulation of rational expression. (Review from Math 940)

Weekly Learning Objectives:

- 1) Use long division to divide polynomials.**
- 2) Use synthetic division to divide polynomials.**
- 3) Use the division algorithm.**
- 4) Use the remainder and factor theorems.**

Dividing Polynomials

Review of Long Division:

$$x-2 \overline{) 12x^3 - 5x^2 - 14x + 1}$$

$$x-1 \overline{) 3x^4 - x^2 + 1}$$

Review of Synthetic Division: (can only be used when dividing by a LINEAR factor!)

$$(12x^3 - 5x^2 - 14x + 1) \div (x - 2)$$

$$x - 1 \overline{) 3x^4 - x^2 + 1}$$

The Division Algorithm summarizes the results of any polynomial division.

If $P(x)$ and $D(x)$ are polynomials, with $D(x)$ not equal to zero, then there exist unique polynomials $Q(x)$ and $R(x)$ such that

$$P(x) = D(x) \cdot Q(x) + R(x)$$

where $R(x)$ is either 0 or of degree less than the degree of $D(x)$.

Show how the Division Algorithm applies to the last example.

Note: If the divisor in the Division Algorithm is of the form $x - c$ for some real number c then the remainder must be a constant. The Division Algorithm could be restated in this situation as:

Let $x = c$ in the above equation:

Remainder Theorem: If the polynomial $P(x)$ is divided by $x - c$, then the remainder is the value $P(c)$.

What is the remainder when dividing $x^{14} - 3x^{12} + x^{10} - 4x + 5$ by $x - 1$?

Use synthetic division to solve: $(x^3 - x^2 + 5x - 2) \div (x - 2)$

What other information do we get because of the Remainder Theorem?

Factor Theorem: $P(c) = 0$ if and only if $x - c$ is a factor of $P(x)$.

Proof:

Is $x-1$ a factor of $x^{12} - 3x^{10} + 5x^8 + 2x^5 - 4x^2 - 1$?

State any theorems you are using.

Reminder: For every zero there is an associated factor.

Find a polynomial function of degree 4 with zeros 3, -2, 0 and 1.