

# The Complex Plane and De Moivre's Theorem Video Lecture

## Section 9.3

### Course Learning Objectives:

Graph polar equations with and without technology.

### Weekly Learning Objectives:

- 1) Plot points in the complex plane.
- 2) Convert a complex number from rectangular form to polar form.
- 3) Find products and quotients of complex numbers in polar form.
- 4) Use De Moivre's theorem.
- 5) Find complex roots.

## The Complex Plane and De Moivre's Theorem

Recall that a complex number is of the form  $x + yi$  where  $x$  and  $y$  are any real numbers.

If we let  $z = x + yi$ , then we can show this geometrically as the point  $(x, y)$  in the  $xy$ -plane.

Using this model, each point in the plane corresponds to a complex number and every complex number corresponds to a point in the plane. We call this the **complex plane**. The  $x$  - axis is the **real axis**, because any point that lies on the real axis is of the form  $z = x + 0i = x$ , a real number. The  $y$  - axis is called the **imaginary axis**, because any point that lies on it is in the form  $z = 0 + yi = yi$ , a pure imaginary number.

Plot the point  $z = 2 - i$  in the complex plane.

Definition: Let  $z = x + yi$  be a complex number. The **magnitude** or **modulus** of  $z$ , denoted by  $|z|$ , is defined as the distance from the origin to the point  $(x, y)$ . That is,

$$|z| = \sqrt{x^2 + y^2}$$

The magnitude of  $z$  is sometimes called the **absolute value** of  $z$ .

Recall that if  $z = x + yi$ , then its conjugate, denoted by  $\bar{z}$ , is  $\bar{z} = x - yi$ .

Notice that  $z\bar{z} =$

Therefore,  $|z| = \sqrt{z\bar{z}}$

When a complex number is written in the standard form  $z = x + yi$ , we say it is in **rectangular**, or **Cartesian form**, because  $(x, y)$  are the rectangular coordinates of the corresponding point in the complex plane. Suppose that  $(r, \theta)$  are the polar coordinates of this point, then

$$x = r\cos\theta \quad y = r\sin\theta$$

Then we can write the complex number  $z = x + yi$  in **polar form** as:

$$z = x + yi = (r \cos\theta) + (r \sin\theta)i = r(\cos\theta + i \sin\theta)$$

The angle  $\theta$ , with  $0 \leq \theta < 2\pi$ , is called the **argument of  $z$** .  
The magnitude of  $z$  is  $|z| = r$ .

Write the expression for  $z = \sqrt{3} - i$  in polar form.

Plot the point corresponding to  $z = 2(\cos 30^\circ + i \sin 30^\circ)$  in the complex plane, and write an expression for  $z$  in rectangular form.

The polar form of a complex number provides an alternative method for finding products and quotients of complex numbers.

**Theorem:** Let  $z_1 = r_1(\cos \theta_1 + i \sin \theta_1)$  and  $z_2 = r_2(\cos \theta_2 + i \sin \theta_2)$  then,

$$z_1 z_2 = r_1 r_2 [\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2)]$$

$$\frac{z_1}{z_2} = \frac{r_1}{r_2} [\cos(\theta_1 - \theta_2) + i \sin(\theta_1 - \theta_2)]$$

Proof:

If  $z = 3(\cos 20^\circ + i \sin 20^\circ)$  and  $w = 5(\cos 100^\circ + i \sin 100^\circ)$ , find the following (leave your answer in polar form):

a)  $zw$

b)  $\frac{z}{w}$

Let  $z = r(\cos \theta + i \sin \theta)$ , then find:

$$z^2 =$$

$$z^3 =$$

$$z^4 =$$

**De Moivre's Theorem:** If  $z = r(\cos \theta + i \sin \theta)$  is a complex number, then

$$z^n = r^n(\cos n\theta + i \sin n\theta)$$

where  $n \geq 1$  is a positive integer.

Find  $[2(\cos 20^\circ + i \sin 20^\circ)]^3$  in standard complex form.

Write  $(1 + i)^5$  in standard form.

Now we will focus on finding roots of complex numbers. If  $w$  is a complex number, then the solutions to  $z^2 = w$  would give the two complex square roots of  $w$ . The solutions to  $z^3 = w$  would give the three complex cube roots of  $w$ , etc.

**If  $w = r(\cos \theta_0 + i \sin \theta_0)$  is a complex number and  $n$  is an integer with  $n \geq 2$ , then there will be  $n$  distinct complex roots of  $w$  given by the formula:**

$$z_k = \sqrt[n]{r} \left[ \cos \left( \frac{\theta_0}{n} + \frac{2k\pi}{n} \right) + i \sin \left( \frac{\theta_0}{n} + \frac{2k\pi}{n} \right) \right]$$

where  $k = 0, 1, 2, 3, \dots, n - 1$ .

Find the complex cube roots of  $-1 + \sqrt{3}i$ . Leave your answer in polar form, with the argument in degrees.